arXiv:math/0211069v1 [math.GT] 4 Nov 2002

UNIVERSAL SPACES FOR ASYMPTOTIC DIMENSION

A. DRANISHNIKOV AND M. ZARICHNYI

Abstract. We construct a universal space for the class of proper metric spaces of bounded geometry and of given asymptotic dimension. As a consequence of this result, we establish coincidence of the asymptotic dimension with the asymptotic inductive dimensions.

1. Introduction

Asymptotic dimension asdim of a metric space was defined by Gromov for studying asymptotic invariants of discrete groups [14]. Then a successful application of the asymptotic dimension was found by G. Yu [19]. He proved the Higher Novikov Signature conjecture for finitely presented groups Γ with a finite asymptotic dimension considered as metric spaces with the word metric. The word metric d_S on a group Γ depends on a generating set $S \subset \Gamma$. The distance $d_S(x, y)$ in the word metric is the minimal length of presentations of the word $x^{-1}y$ in the alphabet S. It turns out that the metric spaces (Γ, d_S) and $(\Gamma, d_{S'})$ are coarsely equivalent (see Section 2 for the exact definition) for any two finite generating sets S and S', and hence $\operatorname{asdim}(\Gamma, d_S) = \operatorname{asdim}(\Gamma, d_{S'})$. Thus, in the case of a finitely generated group Γ, one can speak about its asymptotic dimension $\operatorname{asdim}\Gamma$ without refereeing to a generating set. In view of Yu's theorem, a finite-dimensionality results are of a particular interest. In [14] Gromov proved that all the hyperbolic groups have finite asymptotic dimension. In [11] the finite dimensionality was established for all Coxeter groups. In [2, 3] finite dimensionality theorems were proved for the fundamental group of graphs of groups. In particular, it was shown that the asymptotic finite dimensionality is preserved under the amalgamated product and the HNN-extension. In [3] an upper estimate for asdim of the fundamental group of a graph of groups is given which turns out to be exact in many cases. Strangely, it did not give an exact estimate in the case of the free product which is seemingly the simplest case of the graph of groups. Only recently the exact formula has been established in [4]

1991 *Mathematics Subject Classification.* 54F45.

Key words and phrases. Asymptotic dimension, universal space, asymptotic inductive dimension.

The paper was written during the second author's visit to the University of Florida. He thanks the Department of Mathematics for hospitality.

with the use of the inductive dimension asInd introduced in [10]. The equality asdim = asInd is proven in the present paper by means of universal spaces.

In the classical dimension theory there are many embedding theorems of different kind. Here we mension three:

(1) Nöbeling-Pontryagin Theorem: Every compactum X with $\dim X \leq n$ can be embedded in $\mathbb{R}^{2n+1}$.

(2) Bowers Theorem [6]: Every compactum X with $\dim X \leq n$ can be embedded in the product of $n+1$ dendrites.

(3) Lefschetz-Menger Theorem [5]: For every n there exists a universal Menger compactum μ^n, i.e. every compactum X with $\dim X \leq n$ can be embedded in μ^n.

We are interested in the asymptotic analogs of these theorems. The interest is partly motivated by a topological approach to Yu's theorem taken in [9]. First, we note that in the large scale world there are direct analogs to (1)–(3) for $n = 0$. Here we give a description of a space M^0 analogous to the Cantor set μ^0. We recall that the Cantor set is the set of all numbers $0 \leq x \leq 1$ which satisfy the following property:

(*) *x can be written without use of 1 in the tercimal system (based on $\{0,1,2\}$).*

Then M^0 can be described as the set of all $x \in \mathbb{R}_+$ satisfying (*). Every proper metric space X with $\operatorname{asdim} X = 0$ and of bounded geometry (see the definition below) can be embedded in the coarse sense in M^0 and hence in $\mathbb{R}$.

The direct analogy between classic and asymtotic dimension theories ends for $n > 0$. It is easy to show that the free group F_2 of two generators has dimension $\operatorname{asdim} F_2 = 1$. It cannot be embedded into $\mathbb{R}^N$, since F_2 has an exponential volume growth. So there is no direct asymptotic analogy to (1). An asymptotic analogy of (2) was found in [9]: Every metric space of bounded geometry can be coarsely embedded in the product of $n+1$ locally finite trees. Since the above product of trees can be embedded in $(2n+2)$-dimensional nonpositively curved manifold, it gives some hope for analogy in the case of (1).

The main goal of this paper is an attempt to find a coarse version of the Lefschetz-Menger theorem, i.e. to construct universal spaces for the asymptotic dimension. We obtain a rather partial result (Theorem 3.10): For every n there is a separable metric space M_n with $\operatorname{asdim} M_n = n$ which is universal for proper metric spaces X with bounded geometry and with $\operatorname{asdim} X \leq n$. Our universal space is neither proper nor of bounded geometry.

The construction of the space M_n is analogous to those for the Menger and Nöbeling spaces. Besides the asymptotic strategy, the main difference is that we are building our fractals not in $\mathbb{R}^{2n+1}$ as in the classical case but in the product of trees. If one accomplish the local (classical) construction in the product of (finite) trees or even dendrites he will get the Menger space according to Bestvina's criterion [5]. We construct our spaces M_n out of product of locally finite trees. For locally finite trees this construction works even better but we have problems with proving universality. Anyway, we construct for every proper metric space of bounded geometry X with asdim an embedding $X \hookrightarrow \prod_{i=0}^{n} T_i$ into the product of $n+1$ finite tree in such a way that X lies in the 'n-skeleton' $M(\{T_i\})$ of $\prod_{i=0}^{n} T_i$. This space $M(\{T_i\})$ has the property $\operatorname{asdim} M(\{T_i\}) = n$. Moreover, $\operatorname{asInd} M(\{T_i\}) = n$. This allows to prove the inequality $\operatorname{asInd} X \leq \operatorname{asdim} X$ for proper metric spaces. For every X the constructed space $M(\{T_i\})$ has the Higson property (see Section 4). There were hopes that the mentioned result on large scale embeddings of asymptotically n-dimensional spaces of bounded geometry into products of $n+1$ nonpositively curved surfaces can be strengthened so that one can take the hyperbolic planes as these manifolds. It turns out, however, (see Theorem 3.15) that there is no proper metric space of finite asymptotic dimension into which every asymptotically 1-dimensional space of bounded geometry can be embedded in the large scale sense.

2. PRELIMINARIES

The generic metric will be denoted by d. If A is a subset of a metric space X and $r \in \mathbb{R}$, then $N_r(A)$ is defined as $\{x \in X \mid d(x,A) < r\}$ if $r > 0$ and $A \setminus \{x \in X \mid d(x,A) < -r\}$ otherwise. The closed r-ball centered at $x \in X$ will be denoted as $B_r(x)$.

A map $f\colon X \to Y$ of metric is called (λ, s)-*Lipschitz* if $d(f(x), f(y)) \leq \lambda d(x,y) + s$ for every $x, y \in X$. The $(\lambda, 0)$-Lipschitz maps are also called λ-Lipschitz; the 1-Lipschitz maps are called *Lipschitz* or *short*. An *asymptotically Lipschitz* map is a map which is (λ, s)-Lipschitz for some $\lambda > 0$, $s > 0$.

For a cover $\mathcal{U}$ of a metric space X, we denote by $L(\mathcal{U})$ its Lebesgue number, $L(\mathcal{U}) = \inf\{\sup\{d(x, X \setminus U) \mid U \in \mathcal{U}\} \mid x \in X\}$, and for a family $\mathcal{U}$ of subsets of a metric space we denote by $\operatorname{mesh}(\mathcal{U})$ the lowest upper bound of the diameters of the elements of $\mathcal{U}$. A family $\mathcal{A}$ of subsets of a metric space is called *uniformly bounded* if there exists a number $C > 0$ such that $\operatorname{diam}(A) < C$ for every $A \in \mathcal{A}$.

For $r > 0$, a metric space X is called r-*discrete* if $d(a,b) \geq r$ for every $a, b \in X$, $a \neq b$. A metric space X is called *discrete* if X is r-discrete for some $r > 0$.

For $r > 0$, the r-*capacity* of a subset Y of a metric space is the maximal cardinality $K_r(Y)$ of an r-discrete subset of Y. A metric space X is of *bounded geometry* if there exists a number $r > 0$ and a function $c\colon [0,\infty) \to [0,\infty)$ such that the r-capacity of every ε-ball $B_\varepsilon(x)$ does not exceed $c(\varepsilon)$.

A map $f\colon X \to Y$ between metric spaces is called *uniformly cobounded* if for every $r > 0$ there exists $C > 0$ such that for every $y \in Y$ the diameter of the set $B_r(y)$ does not exceed C.

A space X is said to be a *geodesic metric space* if for every two points $x, y \in X$ there is an isometric embedding $j\colon [0, d(x,y)] \to X$ such that $j(0) = x$ and $j(d(x,y)) = y$.

A geodesic segment that connects points x, y in a geodesic metric space will be denoted by $[x, y]$.

A metric space is called *uniformly arcwise connected* if for every $\varepsilon > 0$ there exists $\delta > 0$ such that for every two points $x, y \in X$ with $d(x, y) < \varepsilon$ there exists a path in X connecting x and y and of diameter $\leq \delta$.

The (finite) products of metric spaces are always endowed with the sup-metric.

Let $\mathcal{C}$ a decomposition of a proper metric space X. Define the *quotient pseudometric* ϱ on X by the following rule: $\varrho(x, y)$ is the greatest lower bound of the sums of the form $\sum_{i=1}^{k-1} d(x_{2i}, x_{2i+1})$, where $x = x_1$, $y = x_{2k}$, and for every $i = 1, \dots, k$ there exists an element $C_i \in \mathcal{C}$ such that $\{x_{2i-1}, x_{2i}\} \subset C_i$. Obviously, the identity map $\mathrm{id}\colon (X, d) \to (X, \varrho)$ is short.

A map $f\colon X \to Y$ between metric spaces is called *coarse uniform* if for every $C > 0$ there is $K > 0$ such that for every $x, x' \in X$ with $d(x, x') < C$ we have $d(f(x), f(x')) < K$. A map $f\colon X \to Y$ is called *metric proper* if the preimage $f^{-1}(B)$ is bounded for every bounded set $B \subset Y$. A map is *coarse* if it is both coarse uniform and metric proper. Two maps $f, g\colon X \to Y$ between metric spaces are called *bornotopic* (see [17]) if there is $C > 0$ such that $d(f(x), g(x)) < C$ for every $x, y \in X$. Two metric space X and Y are *coarse equivalent* if there are coarse maps $f\colon X \to Y$, $g\colon Y \to X$ such that the compositions fg and gf are bornotopic to the corresponding identity maps.

Lemma 2.1. [8] *Let X be a geodesic metric space and $f\colon X \to Y$ be a coarse uniform map. Then f is an asymptotically Lipschitz map.*

We will need the following simple result whose proof mimicks that of Proposition 2 from [10].

Lemma 2.2. *For every proper metric space X and every $C > 0$ there is a C-discrete subset Y of X such that $d(x, Y) \leq C$ for every $x \in X$.*

A map $f\colon X \to Y$ of metric spaces is called a *large scale embedding* (see [17]) if there exist increasing functions $\varphi_1, \varphi_2\colon [0, \infty) \to [0, \infty)$ with $\lim_{t\to\infty} \varphi_1(t) = \lim_{t\to\infty} \varphi_2(t) = \infty$ such that $\varphi_1(d(x, y)) \leq d(f(x), f(y)) \leq \varphi_2(d(x, y))$ for every $x, y \in X$. It is easy to see that every metric space is coarsely equivalent to its image under a coarse embedding.

2.1. **Asymptotic dimension.**

The notion of asymptotic dimension is introduced by Gromov [14].

Definition 2.3. The *asymptotic dimension* of a metric space X does not exceed n (written $\operatorname{asdim} X \leq n$) if for every $D > 0$ there exists a uniformly bounded cover $\mathcal{U}$ of X such that $\mathcal{U} = \mathcal{U}^0 \cup \dots \cup \mathcal{U}^n$, where all $\mathcal{U}^i$ are D-disjoint.

A *piecewise-euclidean* n-dimensional complex of mesh D is a complex which is a union of isometric copies of the standard n-dimensional symplex of mesh D in $\mathbb{R}^{n+1}$.

Lemma 2.4. *For a proper metric space X the following two conditions are equivalent:*

1) $\operatorname{asdim} X \le n$;
2) *for any $D > 0$ there is a uniformly cobounded short proper map of X into a piecewise-euclidean complex of mesh D and dimension n.*

Proof. See [14]. □

A family of metric spaces X_α satisfies the inequality $\operatorname{asdim} X_\alpha \le n$ *uniformly* (see [2]) if for arbitrary large $D > 0$ there exists $R > 0$ and R-bounded D-disjoint families $\mathcal{U}_\alpha^0, \dots \mathcal{U}_\alpha^n$ such that $\cup_{i=0}^n \mathcal{U}_\alpha^i$ is a cover of X_α.

Theorem 2.5. [2] *Assume that $X = \cup_\alpha X_\alpha$ and $\operatorname{asdim} X_\alpha$ uniformly. Suppose that for any R there is $Y_R \subset X$ with $\operatorname{asdim} Y_R \le n$ and such that the family $\{X_\alpha \setminus Y_R\}$ is R-disjoint. Then $\operatorname{asdim} X \le n$.*

3. ASYMPTOTIC EMBEDDINGS INTO PRODUCT OF TREES

3.1. **Trees.** A geodesic metric space T is called an $\mathbb{R}$-*tree* (a real tree) if (1) every two points in T are connected by a unique geodesic segment and (2) if $[x, y] \cap [y, z] = \{y\}$ then $[x, y] \cup [y, z] = [x, z]$ for all $x, y, z \in T$.

Recall that in this definition $[a, b]$ stands for a geodesic segment connecting a and b. We will also use self-explaining notations $[a, b)$ and (a, b) for (half)open geodesic segments.

Every tree (connected connected acyclic graph) is an $\mathbb{R}$-tree. We assume that the graphs are endowed with a geodesic metric whose restriction to every edge is isometric to the standard unit segment.

A (half)open segment in T is *free* if it is an open subset of T.

The *mesh* of T is the greatest lower bound of the diameters of the maximal (with respect to the inclusion) free (half)open segments in T.

Definition 3.1. An $\mathbb{R}$-tree T is said to be *regular* if there exists a sequence of $\mathbb{R}$-trees $T = T_0 \supset T_1 \supset T_2 \supset \dots$ with the following properties:

1) $\cap_{i=0}^\infty T_i = \emptyset$;
2) for every i, the set $T_i \setminus T_{i+1}$ is a disjoint union of a uniformly bounded family of maximal free in T_i half-open segments;
3) $D_0 < D_1 < D_2 < \dots$ and $\lim_{i\to\infty} D_i = \infty$, where D_i denotes the mesh of the $\mathbb{R}$-tree T_i.



If in the above definition $D_i = 2^i$, we say that T is *binary regular*.

Suppose that $T^{(0)}, \dots, T^{(n)}$ is a sequence of regular $\mathbb{R}$-trees. For every j, a sequence of $\mathbb{R}$-trees $T^{(j)} = T_0^{(j)} \supset T_1^{(j)} \supset T_2^{(j)} \supset \dots$ is given such that properties 1)–3) from Definition 3.1 are satisfied with $T_i = T_i^{(j)}$. For every i and j denote by $r_i^{(j)} \colon T^{(j)} \to T_i^{(j)}$ the retraction that maps every component C of the set $T^{(j)} \setminus T_i^{(j)}$ onto its boundary ∂C (which is a singleton). We will refer to these retractions as the canonical retractions. For $k \le i$ we have $r_i^{(j)} = r_{ki}^{(j)} r_k^{(j)}$. Let

$$M_j = \left\{ (x_0, \dots, x_n) \in \prod_{i=0}^{n} T^{(i)} \mid \text{there exists } i \text{ such that } r_i(x_i) \in \partial(T_{j-1}^{(i)} \setminus T_j^{(i)}) \right\}$$

and $M = M(T^{(0)}, \dots, T^{(n)}) = \cap_{j=1}^{\infty} M_j$.

Denote by $\mathcal{M}_n$ the class of spaces of the form $M(T^{(0)}, \dots, T^{(n)})$ for some sequence $T^{(0)}, \dots, T^{(n)}$ of regular $\mathbb{R}$-trees.

Suppose that T, S are regular $\mathbb{R}$-trees given with filtrations $(T_j)_{j=0}^{\infty}$, $(S_j)_{j=0}^{\infty}$ satisfying the conditions from Definition 3.1. We denote by $r_j \colon T \to T_j$, $\varrho_j \colon S \to S_j$ the canonical retractions.

Definition 3.2. A map $f \colon T \to S$ is said to be *regular* if the following conditions are satisfied:

(i) $f(T_j) \subset S_j$ for every j;
(ii) $f(r_j(\partial(T_{j-1} \setminus T_j)) \subset \varrho_j(\partial(S_{j-1} \setminus S_j))$ for every j.

The construction $M(T^{(0)}, \dots, T^{(n)})$ is functorial in the following sense. Suppose that T^i, S^i are regular $\mathbb{R}$-trees given with filtrations $(T^i_j)_{j=0}^{\infty}$, $(S^i_j)_{j=0}^{\infty}$, $i \in \{0, 1, \dots, n\}$ satisfying the conditions from Definition 3.1. We denote by $r^i_j \colon T^i \to T^i_j$, $\varrho^i_j \colon S^i \to S^i_j$ the canonical retractions. Suppose that $f_i \colon T^i \to S^i$ are regular maps.

Lemma 3.3. *Under these conditions*

$$(f_0 \times \cdots \times f_n)(M(T^{(0)}, \dots, T^{(n)})) \subset M(S^{(0)}, \dots, S^{(n)}).$$

Proof. Obvious. □

Proposition 3.4. *For every* $M \in \mathcal{M}_n$ *we have* $\operatorname{asdim} M \le n$.

Proof. We apply Lemma 2.4. Note that for every j the map

$$r_j = \prod_{i=0}^{n} r_j^{(i)} \colon \prod_{i=0}^{n} T^{(i)} \to \prod_{i=0}^{n} T^{(i)}$$

maps M onto an n-dimensional piecewise-euclidean polyhedron of mesh D_j. Since r_j is uniformly cobounded, we are done. □

Theorem 3.5. *For every proper metric space* X *of* $\operatorname{asdim} X \le n$ *there exist locally finite binary regular trees* $T^0, \dots, T^n$ *such that* X *is large scale embeddable into* $M(T^0, \dots, T^n)$.

The proof is a modification of the corresponding result from [8].

Proposition 3.6. *Let* X *be a proper metric space with base point* x_0 *and* $\operatorname{asdim} X \le n$. *Then there exists a sequence* $(\mathcal{U}_k)_{k=1}^{\infty}$ *of open covers of* X *such that every* $\mathcal{U}_k$ *splits into the union* $\mathcal{U}_k = \mathcal{U}_k^0 \cup \dots \cup \mathcal{U}_k^n$ *of* d_k*-disjoint families and the following conditions are satisfied:*

1) $L(U_k) > d_k$ *and for every* i *and every* $U \in \mathcal{U}_k^i$ *we have* $N_{-d_k}(U) \ne \emptyset$;
2) $d_{k+1} = 2^{2k+2} m_k$, *where* m_k *is the mesh of* $\mathcal{U}_k$;
3) *for every* $m \in \mathbb{N}$ *and for every* i *there is* $k \in \mathbb{N}$ *and an element* $U \in \mathcal{U}_k^i$ *such that* $N_m(x_0) \subset U$.
4) *for every* $k, l,\ k < l$ *and every* $U \in \mathcal{U}_k^i$, $V \in U_l^i$, *if* $U \not\subset V$, *then* $d(U, V) \ge d_k/2$.

Proof. We start with repeating the construction from the proof of Proposition 1 from [9]. We proceed by induction. Let $\hat{\mathcal{U}}_0$ be an open cover of X which is $\hat{d}_0$-discrete with $\hat{d}_0 > 2$. We enumerate the partition $\hat{\mathcal{U}}_0 = \hat{\mathcal{U}}_0^0 \cup \dots \cup \hat{\mathcal{U}}_0^n$ in such a way that $d(x_0, X \setminus U) > \hat{d}_0$ for some $U \in \hat{U}_0^0$.

Assume that families $\hat{\mathcal{U}}_k$ are constructed for all $k \le l$. Define $d_{l+1} = 2^{l+2} m_l$ and consider a uniformly bounded cover $\bar{\mathcal{U}}_{l+1}$ with $L(\bar{\mathcal{U}}_{l+1}) > 2d_{l+1}$ and such that there is a splitting $\bar{\mathcal{U}}_{l+1} = \bar{\mathcal{U}}_{l+1}^0 \cup \dots \cup \bar{\mathcal{U}}_{l+1}^n$, where $\bar{\mathcal{U}}_{l+1}^i$ are d_{l+1}-disjoint and $N_{-2d_{l+1}}(U) \ne \emptyset$ for all $U \in \bar{\mathcal{U}}_{l+1}$. The families $\bar{\mathcal{U}}_{l+1}^0, \dots, \bar{\mathcal{U}}_{l+1}^n$ are enumerated in such a way that $d(x_0, X \setminus U) > 2d_{l+1}$ for some $U \in \bar{\mathcal{U}}_{l+1}^i$, where $i = l + 1 \mod n + 1$. For every $U \in \bar{\mathcal{U}}_{l+1}^i$ let

$$\tilde{U} = U \setminus \{\overline{N_4(V)} \mid V \in \mathcal{U}_k^i,\ k \le l,\ V \not\subset U\}$$

and $\mathcal{U}_{l+1}^i = \{\tilde{U} \mid U \in \bar{\mathcal{U}}_{l+1}^i\}$.

It is proved in [9] that the sequence $(\hat{\mathcal{U}}_k)$ satisfies the following properties:

1) $L(\hat{\mathcal{U}}_k) > d_k$ and for every i and every $U \in \hat{\mathcal{U}}^i_k$ we have $N_{-d_k}(U) \neq \emptyset$;
2) $d_{k+1} = 2^k m_k$, where m_k is the mesh of $\hat{\mathcal{U}}_k$;
3) for every $m \in \mathbb{N}$ and for every i there is $k \in \mathbb{N}$ and an element $U \in \hat{\mathcal{U}}^i_k$ such that $N_m(x_0) \subset U$;
4) for every $U \in \hat{\mathcal{U}}^i_k$, $V \in \hat{\mathcal{U}}^i_l$, $U \not\subset V$ we have $d(U,V) \geq 4$.

We are going to modify the sequence $(\hat{\mathcal{U}}_k)_{k=1}^\infty$. Passing, if necessary, to a subsequence of the sequence $(\hat{\mathcal{U}}_k)_{k=1}^\infty$ we may assume that the sequence $(\hat{\mathcal{U}}_k)_{k=1}^\infty$ itself satisfies conditions 1), 3),4) and the following condition:

2*) $d_{k+1} \geq 2^{4k+6} m_k$.

Define, by induction, numbers $\tilde{d}_k$ and families $\tilde{\mathcal{U}}^i_k$. Let $\tilde{d}_0 = d_0$ and $\tilde{\mathcal{U}}_0 = \mathcal{U}_0$ and suppose $\tilde{d}_j$ and $\tilde{\mathcal{U}}_j$ have been defined for every $j < k$. For every $U \in \mathcal{U}^i_k$ apply induction by p to define the sets $\tilde{U}(p)$. Let $\tilde{\mathcal{U}}(0) = U$ and suppose sets $\tilde{U}(p)$ have been defined for all $p < q$, for some $q \leq k$. Let

$$\tilde{\mathcal{U}}(q) = \tilde{U}(q-1) \cup \bigcup \{V \in \tilde{\mathcal{U}}^i_q \mid d(\tilde{U}(q-1), V) < \tilde{d}_q/2\}.$$

Then, by the definition, $\tilde{U} = \tilde{U}(k-1)$ and $\tilde{U}^i_k = \{\tilde{U} \mid U \in \mathcal{U}^i_k\}$.

We are going to verify conditions 1)–4) (with U_k, d_k, and m_k replaced by $\tilde{U}_k$, $\tilde{d}_k$, and $\tilde{m}_k$ respectively). Conditions 1) and 3) are obvious.

Denote by $\tilde{m}_k$ the mesh of $\tilde{\mathcal{U}}^i_k$ and let

$$\tilde{d}_k = d_k - 2^{k+2}\tilde{m}_{k-1}. \tag{3.1}$$

Note that for every $\tilde{U}, \tilde{V} \in \tilde{U}^i_k$, $\tilde{U} \neq \tilde{V}$ we have

$$\begin{aligned}
d(\tilde{U}, \tilde{V}) =& d(\tilde{U}(k-1), \tilde{V}(k-1)) \geq d(\tilde{U}(k-2), \tilde{V}(k-2)) - \tilde{d}_{k-1} - 2\tilde{m}_{k-1} \\
\geq& d(\tilde{U}(k-2), \tilde{V}(k-2)) - 3\tilde{m}_{k-1} \\
\geq& d(\tilde{U}(k-3), \tilde{V}(k-3)) - 3\tilde{m}_{k-1} - 3\tilde{m}_{k-2} \\
& \dots \\
\geq& d(U,V) - 3\sum_{i=0}^{k-1} \tilde{m}_i \geq d_k - 3k\tilde{m}_{k-1} \geq \tilde{d}_k,
\end{aligned}$$

and therefore the family $\tilde{\mathcal{U}}^i_k$ is $\tilde{d}_k$-disjoint.

Condition 2). Let $\tilde{U} \in \tilde{\mathcal{U}}^i_k$, then $\tilde{U} = \tilde{U}(k-1)$ and

$$\begin{aligned}
\operatorname{diam} \tilde{U} \leq& \operatorname{diam} \tilde{U}(k-2) + 2\tilde{m}_{k-1} + \tilde{d}_{k-1} \leq \operatorname{diam} \tilde{U}(k-2) + 3\tilde{m}_{k-1} \\
& \dots\dots \\
\leq& \operatorname{diam} \tilde{U}(0) + 3\sum_{j=0}^{k-1} \tilde{m}_j \leq m_k + 3k\tilde{m}_{k-1} \leq m_k + 2^{k+2}\tilde{m}_{k-1}
\end{aligned}$$

(here we used an obvious equality $m_j \geq d_j$, for every j), whence

$$\tilde{m}_k \leq m_k + 2^{k+2}\tilde{m}_{k-1}. \tag{3.2}$$

Then

$$\tilde{m}_k \leq m_k \sum_{j=0}^{k-1} 2^{j(k+2)} \leq m_k 2^{2k+3}$$

and using condition 2^*) we obtain

$$\begin{aligned} \tilde{d}_k =& d_k - 2^{k+2}\tilde{m}_{k-1} \geq d_k - 2^{2k+3}m_{k-1} \\ \geq& 2^{4k+6}m_{k-1} - 2^{2k+3}m_{k-1} \geq 2^{4k+5}m_{k-1} \\ \geq& 2^{2k+2}\tilde{m}_{k-1}. \end{aligned} \tag{3.3}$$

Let us verify condition 4). Suppose the contrary. Then there exist $\tilde{U} \in \tilde{\mathcal{U}}^i_k$, $\tilde{V} \in \tilde{\mathcal{U}}^i_l$, $\tilde{U} \not\subset \tilde{V}$, and $k \leq l$ such that $d(\tilde{U}, \tilde{V}) < d_k/2$. We also suppose that l is the minimal number with that property. Since $d(\tilde{U}, \tilde{V}) < d_k/2$, there exists $x \in \tilde{V}$ such that $d(x, \tilde{U}) < d_k/2$. Let $j \geq 0$ be the minimal integer with the property $x \in \tilde{V}(j)$. It follows from the definition of $\tilde{\mathcal{U}}^i_l$ that $j \leq l-1$. Since $x \in \tilde{V}(j) \setminus \tilde{V}(j-1)$, it follows from the definition of $\tilde{V}(j)$ that there exists $\tilde{W} \in \tilde{\mathcal{U}}^i_j$ such that $x \in \tilde{W}$. Since $j < l$, this contradicts to the choice of l.

□

For every element $U \in \mathcal{U}_i$ denote by $\psi(U)$ the minimal (with respect to inclusion) $V \in \mathcal{U}_i$ such that U is a proper subset of $\psi(U)$.

Proposition 3.7. *Let X be a proper metric space and $(\mathcal{U}_k)$ a sequence of open covers of X satisfying properties 1)–4) from Proposition 3.6. Let $U \in \mathcal{U}_k$ and ϱ be the quotient metric on U with respect to the decomposition of U into singletons and the elements of the family $\mathcal{V} = \psi^{-1}(U)$. For every $V \in \mathcal{V} \cap \mathcal{U}^i_j$, $W \in \mathcal{V} \cap \mathcal{U}^i_l$, where $j \leq l$, we have $\varrho(V, W) \geq 2^{2j}$.*

Proof. The assertion is obvious for $j = 0$. Let $V \in \mathcal{V} \cap \mathcal{U}^i_j$, $W \in \mathcal{V} \cap \mathcal{U}^i_l$, where $1 \leq j \leq l$. Then $\varrho(V, W)$ is the greatest lower bound of the sums of the form $\sum_{i=1}^{p-1} d(x_{2i}, x_{2i+1})$, where $x \in V$, $x_{2p} \in W$, and for every $i = 1, \dots, p$ there exists an element $C_i \in \psi^{-1}(U)$ such that $\{x_{2i-1}, x_{2i}\} \subset C_i$. Without loss of generality, we may assume that in the sums above $C_i \in \psi^{-1}(U) \setminus \mathcal{V}_j$ for all $j < p$. Then $(p-2)m_{j-1} + (p-1)d_{j-1} \geq d_j$ and therefore, by condition 2) of Proposition 3.6,

$$p - 1 \geq \frac{d_j - d_{j-1}}{m_{j-1} + d_{j-1}} \geq \frac{d_j}{4m_{j-1}} \geq 2^{2j}$$

and $\sum_{i=1}^{p-1} d(x_{2i}, x_{2i+1}) \geq (p-1)d_0 \geq 2^{2j}$. Hence $\varrho(V, W) \geq 2^{2j}$. □

Proposition 3.8. *Let X, U, and ϱ be as in Proposition 3.7. There exists $x \in U$ with $\varrho(x, X \setminus U) \geq 2^{2k}$.*

Proof. By condition 1) of Proposition 3.6, there exists $x \in U$ with $d(x, \partial U) \geq d_k$. Arguing like in the proof of Proposition 3.7 we conclude that $\varrho(x, X \setminus U) \geq 2^{2k}$. □

Fix a sequence $(\mathcal{U}_k)$ of open covers of X satisfying properties 1)–4) from Proposition 3.6. Given $U \in \mathcal{U}_k^i$ define a Lipschitz map $f_U \colon \bar{U} \to I_U = [0, 2^{2^k}]$ by the following procedure. Let ϱ denote the pseudometric generated by the decomposition of U into singletons and the elements of the family $\psi^{-1}(U)$ and let $\mathcal{V}_j = (\{V \in \mathcal{U}_l^i \mid j \leq l < k\}) \cap \psi^{-1}(U)$.

We are going to define by induction with respect to j maps $\theta_j, \bar{\theta}_j \colon \partial U \cup \overline{\bigcup \mathcal{V}_j} \to I_U$. Define the map $\theta_1 \colon \partial U \cup \overline{\bigcup \mathcal{V}_{k-1}} \to I_U$ by the formula $\theta_1(x) = 2^{-k}\varrho(x, X \setminus U)$. Let $\bar{\theta}_1(x) = 2^{k-1}[2^{-k+1}\theta_1(x)]$. The map $\bar{\theta}_1$ is a 2^{-k+1}-Lipschitz map into I_U defined on the closed subset $\partial U \cup \overline{\cup \mathcal{V}_{k-1}}$ of $\bar{U}$.

Suppose the maps $\theta_j, \bar{\theta}_j$ are defined for all j, $j_0 < j < k$ and the maps $\bar{\theta}_j$ are 2^{-j}-Lipschitz. By the theorem on extension of Lipschitz maps (see [8]) there exists a 2^{-j_0+1}-Lipschitz extension of $\bar{\theta}_{j_0-1}$ onto the set $\partial U \cup \overline{\bigcup \mathcal{V}_{j_0}}$. Let $\bar{\theta}_{i_0}(x) = 2^{i_0}[2^{-i_0}\theta_{i_0}(x)]$.

As a final result, we obtain a 1-Lipschitz map

$$\bar{\theta}_0 \colon \partial U \cup \overline{\bigcup \mathcal{V}_0} = \partial U \cup \overline{\bigcup \psi^{-1}(U)} \to I_U.$$

Denote by f_U its 1-Lipschitz extension onto $\bar{U}$.

We will follow [9] and construct a locally finite tree T^i by the following procedure. Assign to every $U \in \mathcal{U}_k^i$, $k = 0, 1, \ldots$, a line segment $I_U = [0, 2^k]$. The tree T^i will be the quotient space of the disjoint union $\sqcup\{I_U \mid U \in \cup_{k=0}^{\infty}\mathcal{U}_k^i\}$ with respect to the following equivalence relation $\sim$: suppose $U \in \mathcal{U}_k^i$, $V \in \mathcal{U}_l^i$, where $k < l$ and $U \in \psi^{-1}(V)$.Then $0 \in I_U$ is identified with $f_V(\partial U) \in I_V$. We argue like in the proof of Theorem 3 from [9] to show that the quotient space $\sqcup\{I_U \mid U \in \cup_{k=0}^{\infty}\mathcal{U}_k^i\}/\sim$ is a tree. We are going to show that T^I is locally finite. Assume the contrary and let y be a vertex of infinite order in T^i. Then there exists an infinite family $\mathcal{V} \subset \mathcal{U}^i$ such that $y = q(0_U)$ for every $U \in \mathcal{V}$. Since the set $\psi^{-1}(U)$ is finite for every $U \in \mathcal{U}$, we conclude that there exists an increasing sequence $(k(j))_{j=1}^{\infty}$ of positive integers and sets $U_{k(j)} \in \mathcal{U}_{k(j)}^i$ such that $U_{k(j)} \in \psi^{-1}\mathcal{U}_{k(j+1)}$ and $f_{U_{k(j+1)}}(U_{k(j)}) = 0$. By conditions 3) and 4), there exists $j > 1$ such that $x_0 \in U_{k(j)}$ and $U_{k(1)} \subset U_{k(j)}$. Then, by condition 3), $U_{k(j+1)} \supset N_{m_{k(j)}+2^{2k(j)+2}d_k}(x_0)$ and, since $\operatorname{diam}(U_{k(j)}) \leq m_{k(j)}$, by condition 3), $d(U_{k(j)}, X \setminus U_{k(j+1)}) \geq 2^{2k(j)+2}d_k$ and, therefore, by Proposition 3.8, $\varrho(U_{k(j)}, X \setminus U_{k(j+1)}) \geq 2^{k(j)}$. Hence $f_{U_{k(j+1)}}(U_{k(j)}) \geq 1 \neq 0$ and we obtain a contradiction.

Let $q\colon \sqcup\{I_U \mid U \in \cup_{k=0}^{\infty}\mathcal{U}_k^i\} \to T^i$ be the quotient map. Define the map $f = f^i \colon X \to T^i$ as follows. Let $x \in X$. By condition 3) of Proposition 3.6, there exists minimal k such that $x \in U$ for some $U \in \mathcal{U}^i_k$. Put $f^i(x) = qf_U(x)$. It can be easily seen that f^i is well-defined and is Lipschitz.

The diagonal map $f = (f^i)_{i=0}^n \colon X \to \prod_{i=0}^n T^i$ is Lipschitz. Show that f is a large-scale embedding. Assuming the contrary we can find sequences (x_l) and (y_l) in X such that $d(x_l, y_l) \to \infty$ while $d(f(x_l), f(y_l)) < C$ for some $C > 0$. Let k be a positive integer such that $2^k > C$. There exists an integer l such that $d(x, y) > m_k$. There exists $i \in \{0, \dots, n\}$ such that $x_l \in U$ and $d(x, X \setminus U > d_k$ for some $U \in \mathcal{U}^i_k$. Then $f_U(x) = 2^k$, By Proposition 3.8. Since $d(x, y) > m_k$, we see that $y \notin U$ and $f_U(y) = 0$. Thus, $d(f(x), f(y)) \geq 2^k > C$, which gives a contradiction.

Note that every T^i is a regular tree. Indeed, for every j denote by T^i_j the subspace $q(\sqcup\{I_U \mid U \in \cup_{k=j}^\infty \mathcal{U}^i_k\})$. Let us verify conditions 1)–3) of Definition 3.1 (with T and T_j replaced by T^i and T^i_j respectively). Condition 1) is obvious. The set $T^i_{j+1} \setminus T^i_j$ is a disjoint union of isometric copies of the half-open segment $(0, 2^{j+1}]$ and is therefore uniformly bounded. To verify condition 3) note that the mesh of the tree T^i_j is 2^j, by the construction.

The trees $T^0, \dots, T^n$ together with filtrations $T^i_0 \supset T^i_1 \supset T^i_2 \supset \dots$ determine the subspace $M(T^0, \dots, T^n)$ of $\prod_{i=1}^n T^i$. We are going to show that $f(X) \subset M(T^0, \dots, T^n)$. Suppose $x \in X$. For every $k = 0, 1, \dots$ there is $i(k) \in \{0, 1, \dots, n\}$ such that $x \in U$ for some $U \in \mathcal{U}^{i(k)}_k$. Then $r^{i(k)}_{k+1}(f(x)) \in \partial T^{i(k)}_{k+1} \setminus T^{i(k)}_k$, i. e. $f(x) \in M_{k+1}$. Therefore, $f(x) \in M_{k+1}$ for every k, i. e. $f(x) \in M(T^0, \dots, T^n)$.

3.2. **Universal space.** A class of metric spaces $\mathcal{C}$ is said to be universal for a class of metric spaces $\mathcal{D}$ if for every $D \in \mathcal{D}$ there is a large scale embedding of C into some $C \in \mathcal{C}$.

Theorem 3.5 can be reformulated as follows.

Theorem 3.9. *The class $\mathcal{M}_n$ is universal for the class of proper metric spaces of asymptotic dimension $\leq n$.*

A metric space X is said to be *universal* for a class of metric spaces $\mathcal{D}$ if for every $D \in \mathcal{D}$ there is a large scale embedding of D into X.

Construct an $\mathbb{R}$-tree T by the following procedure. Define inductively a sequence (T_i) of $\mathbb{R}$-trees and retractions $r_i \colon T_i \to T_0$. Let $T_0 = \mathbb{R}_+$. Denote by T_1 an $\mathbb{R}$-tree obtained by attaching to every integer point in $\mathbb{R}_+$ a countable set of isometric copies of the unit segment $[0, 1]$. Denote by r_1 a retraction of T_1 onto T_0 that sends every component C which is constant on every component of the complement $T_1 \setminus T_0$. Suppose that $\mathbb{R}$-trees T_i and retractions r_i are defined for all $i < j$. Denote by S the subtree $r_{j-1}^{-1}([0, 2^j])$ with base point 2^j. The $\mathbb{R}$-tree T_j is obtained from T_{j-1} by attaching to every point of the form $k2^j$, $k \in \mathbb{N}$, of $\mathbb{R}_+ \subset T_{j-1}$ a countable family of isometric copies of S by the base point.

Denote by r_j a retraction of T_j onto T_0 that sends every component C which is constant on every component of the complement $T_j \setminus T_0$.

Let $T = \cup_{i=1}^{\infty} T_i$. It is easy to see that T is a regular $\mathbb{R}$-tree that contains every locally finite binary regular tree so that the inclusion preserves the filtration. By Lemma 3.3, we conclude that for every sequence $T^0, T^1, \dots, T^n$ of locally finite binary regular trees we have

$$M(T^0, T^1, \dots, T^n) \subset M_n = M(S^0, S^1, \dots, S^n),$$

where S^i is isomorphic to T for every $i = 0, 1, \dots, n$.

We therefore obtain the following

Theorem 3.10. *There exists a separable metric space* M_n *with* $\operatorname{asdim} M_n = n$ *universal for the class of proper metric spaces* X *with* $\operatorname{asdim} X \le n$.

For the spaces of asymptotic dimension zero the result can be improved.

Theorem 3.11. *There exists a proper metric space* M_0 *of bounded geometry with* $\operatorname{asdim} M_0 = 0$ *universal for the class of proper metric spaces* X *of bounded geometry with* $\operatorname{asdim} X \le 0$.

Proof. Let M_0 denote the set of integers whoce tercimal expansion consists only from 0s and 1s. Obviously, $\operatorname{asdim} M_0 = 0$ and M_0 is of bounded geometry. For every $k \in \mathbb{N}$ the set M_0 can be represented as the union of 3^k-discrete family of the intervals of length 3^k in the set of natural numbers.

Let X be a proper metric space of bounded geometry with $\operatorname{asdim} X \le 0$. Let $x_0 \in X$ be a base point. There exists a sequence $(\mathcal{U}_k)$ of uniformly bounded covers of X with the following properties:

(1) $\mathcal{U}_k$ refines U_{k+1};
(2) $\mathcal{U}_k$ is k-discrete;
(3) $\cup\{U \in \cup_k \mathcal{U}_k \mid x_0 \in U\} = X$.

For $U \in \mathcal{U}_k$, denote by $\psi(U)$ the unique $V \in \mathcal{U}_{k+1}$ that contains U.

Since X is of bounded geometry, for every $k \in \mathbb{N}$ there exists $C_k > 0$ such that $|\psi^{-1}(U)| \le C_k$ for every $U \in \mathcal{U}_{k+1}$.

Using the mentioned decomposition of M_0 into the union of intervals, we can easily construct a sequence of covers $(\mathcal{V}_k)$ satisfying the properties:

(4) $\mathcal{V}_k$ refines V_{k+1};
(5) $\mathcal{V}_k$ is k-discrete;
(6) $|\psi^{-1}(V)| > C_k$ for every $V \in \mathcal{V}_{k+1}$ (as above, for every $W \in \mathcal{V}_k$ by $\psi(V)$ we denote the unique element of $\mathcal{V}_{k+1}$ that contains V).

For every $U \in \mathcal{U}_1$ and $V \in \mathcal{V}_1$ denote by g_{UV} arbitrary constant map from U to V. Suppose that for every $i \le k$ and every $U \in \mathcal{U}_i$ and $V \in \mathcal{V}_i$ a map $g_{UV} \colon U \to V$ is defined. Given $U \in \mathcal{U}_{k+1}$, $V \in V_{k+1}$, consider arbitrary injective map $\alpha \colon \psi^{-1}(U) \to \psi^{-1}(V)$. Define $g_{UV} \colon U \to V$ as follows: $g_{UV}|W = g_{W\alpha(W)}$ for every $W \in \psi^{-1}(U)$.

Now we are ready to define a map $f\colon X \to M_0$. For every $k \in \mathbb{N}$ denote by U_k the unique element of $\mathcal{U}_k$ that contains x_0. By induction, define a sequence of maps $f_k\colon U_k \to M_0$ with the following properties:

(1′) $f_{k+1}|U_k = f|k$, for every k;

(2′) $f(U_k)$ is contained in an element of $\mathcal{U}_{k+1}$.

Let $f_1 = g_{U_1 V}\colon U_1 \to V \subset M_0$, for some $V \in \mathcal{V}_1$. Suppose that f_i are defined for every $i \le k$. Let $V' \in \mathcal{V}_k$ be such that $f_k(U_k) \subset V'$. Denote by $\alpha\colon \psi^{-1}(\psi(U_k)) \to \psi^{-1}(\psi(V'))$ an embedding such that $\alpha(U_k) = V'$. Define f_{k+1} by the conditions $f_{k+1}|U_k = f_k$ and $f_{k+1}|W = g_{W\alpha(W)}$ for every $W \in \psi^{-1}(\psi(U_k)) \setminus \{U_k\}$.

By condition (1′), the sequence of maps (f_k) uniquely determines a map $f\colon X \to M_0$. Using properties (1)–(6) it is easy to see that f is a coarse embedding. □

3.3. **Nonexistence results.** There is no counterpart of Theorem 3.11 in higher dimensions.

Theorem 3.12. *There is no proper metric space universal for the class of proper metric spaces Y with* $\operatorname{asdim} Y \le 1$.

Proof. Suppose the contrary and let X be a proper metric space universal for the class of proper metric spaces Y with $\operatorname{asdim} Y \le 1$. Let $x_0 \in X$ be a base point. Denote by $\alpha(r)$ the 1-capacity of the ball $N_r(x_0)$, i.e. the number $K_1(N_r(x_0))$. Note that for every $x \in X$ we have

$$K_1(N_r(x)) \le K_1(N_{r+d(x,x_0)}(x_0)) \le \alpha(r + d(x, x_0)).$$

Let T be a locally finite tree with a base point t_0 and the index function $\varphi(r)$ which we specify later. Suppose that $f\colon T \to X$ is a coarse embedding. Since T is a geodesic space, the map f is proper and (λ, s)-Lipschitz for some $\lambda > 0$, $s \ge 0$. There exists $a > 0$ such that $d(x, y) \ge a$ implies $d(f(x), f(y)) \ge 1$. There exist $\lambda', s' > 0$ such that $f(N_r(t_0)) \subset N_{\lambda' r + s'}(x_0)$. Therefore

$$K_a(N_r(t_0) \le K_1(N_{\lambda' r+s'}(x_0)) = \alpha(\lambda' r + s').$$

Now suppose that $\varphi\colon \mathbb{N} \to \mathbb{N}$ satisfies the following property: for every $n \in \mathbb{N}$ there exists $m \in \mathbb{N}$ with $\alpha(nr + n) < \varphi(r)$ for all $r \ge m$. Define T as follows Let $T_0 = \mathbb{R}_+$. For every $m \in \mathbb{N}$ denote by T_m a tree obtained by attaching to every integer point $n \in \mathbb{R}_+$, $n \ge m$, $\varphi(n)$ copies of the segment $[0, m]$ by its endpoints. Let $T = \cup\{T_n \mid n \in \mathbb{N} \cup \{0\}\}$.

Now

$$K_a(N_r(t_0)) \le \alpha(\lambda' r + s') \le \alpha(nr + n),$$

where $n \in \mathbb{N}$, $n \ge \max\{\lambda', s'\}$. Let $m \in \mathbb{N}$, $m \ge a$. Then for every $r \ge 2m + 1$ we obtain

$$K_a(N_r(t_0)) \ge K_m(N_r(t_0)) \ge \varphi([r]),$$

a contradiction with the choice of φ. □

Show that there is no universal proper metric space of given asymptotic dimension in the class of proper metric spaces of bounded geometry.

We fix natural $n \geq 1$.

For natural k, denote by $X(1,k)$ the set

$$\{(x_0,\dots,x_n) \in [0,k]^{n+1} \mid x_i \notin \mathbb{Z} \text{ for at most one } i = 0,\dots,n\}.$$

For natural m, let $X(m,k) = \{mx \mid x \in X(1,k)\}$.

Denote by $\mathbf{V}$ the set of covers $\mathcal{V}$ of $X(1,k)$ such that $\mathcal{V} = \cup_{i=0}^{n}\mathcal{V}^i$, where $\mathcal{V}_i$ are 3-discrete, $i = 0,1,\dots,n$. For $\mathcal{V} \in \mathbf{V}$, denote by $\mu_{\mathcal{V}}(k)$ the maximal 1-capacity of $V \in \mathcal{V}$ and let $\mu_k = \min\{\mu_{\mathcal{V}}(k) \mid \mathcal{V} \in \mathbf{V}\}$.

Given $\lambda > 0$, we say that a subset A of a metric space X is a $< \lambda$-*component* of X if A is a maximal (with respect to inclusion) subset of B with respect to the property that every two points of A can be connected by a λ'-chain, for some $\lambda' < \lambda$.

Lemma 3.13. *$\mu_k \to \infty$ as $k \to \infty$.*

Proof. Assume the opposite, i.e. there exists a constant S such that $\mu_k \leq S$, for all k. Given $\mathcal{V} \in \mathbf{V}$ with $\mathcal{V}_{i=0}^{\infty} = \cup\mathcal{V}^i$, where $\mathcal{V}^i$ are 3-discrete, $i = 0,\dots,n$, for every $V \in \mathcal{V}$, denote by $\tilde{V}$ the family of $<$ 3-connected components of V. Let $\tilde{\mathcal{V}}_i = \cup\{\tilde{V} \mid V \in \mathcal{V}_i\}$, $i = 0,\dots,n$, and $\tilde{\mathcal{V}} = \cup_{i=0}^{n}\tilde{\mathcal{V}}_i$. From our assumption it easily follows that there exists a constant $C > 0$ such that for every $W \in \tilde{\mathcal{V}}$ we have $\operatorname{diam}(W) \leq C$.

Now, for every $W \in \tilde{\mathcal{V}}$ define $\hat{W}$ as the 1-neighborhood of W in $[0,k]^{n+1}$. The families $\{\hat{W} \mid W \in \tilde{\mathcal{V}}_i\}$, $i = 0,1,\dots,n$, are 1-discrete, of mesh $\leq C$, and together form a cover of $[0,k]^{n+1}$. Then the families $\{(1/k)\hat{W} \mid W \in \tilde{\mathcal{V}}_i\}$, $i = 0,1,\dots,n$, are discrete, of mesh $\leq C/k$, and together form a cover of $[0,1]^{n+1}$. Because of arbitrarity of k, we obtain a contradiction with Ostrand's characterization of the covering dimension [16]. $\square$

Passing to the images of spaces under the homothety with coefficient m and centered at the origin, we derive from Lemma 3.13 the following statement.

Proposition 3.14. *For every $m, s \in \mathbb{N}$ there exists $r(m,s) \in \mathbb{N}$ such that for any $r \geq r(m,s)$ the following holds: given a cover $\mathcal{V}$ of $X(m,r)$ that splits into union of $n+1$ $3m$-disjoint families, there is $V \in \mathcal{V}$ with $K_m(V) \geq s$.*

Theorem 3.15. *There is no proper metric space Y of bounded geometry,* $\operatorname{asdim} Y = n$*, with the following property: for every proper metric space X of bounded geometry,* $\operatorname{asdim} X = n$*, there exists a large scale embedding of X into Y.*

Proof. Suppose the opposite and let Y be such a space. There exists a sequence of covers $(\mathcal{U}_k)$ of Y with the following properties:

1) $\mathcal{U}_k$ is uniformly bounded;

2) $\mathcal{U}_k = \cup_{i=0}^n \mathcal{U}_k^i$, where $\mathcal{U}_k^i$ are d_k-discrete with $d_k \ge k^2$.

Since Y is of bounded geometry, $c_k = \max\{K_1(U) \mid U \in \mathcal{U}_k\} < \infty$.

For every $m \in \mathbb{N}$, choose R_m so that for any $r \ge r(m, R_m)$ the following holds: given a cover $\mathcal{V}$ of $X(m,r)$ that splits into union of two $3m$-disjoint families, there is $V \in \mathcal{V}$ with $K_m(V) \ge \sum_{i=1}^m c_m + 1$.

Define a space X as follows: attach to $\mathbb{R}_+$ at every point $m \in \mathbb{N} \subset \mathbb{R}_+$ a copy of space $X(m, R_m)$, with the geodesic metric on the adjunction space. Note that $\operatorname{asdim} X = 1$.

Suppose that $f\colon X \to Y$ is a large scale embedding, i. e. there exist monotone, increasing to infinity functions $\varphi_1, \varphi_2\colon \mathbb{R}_+ \to \mathbb{R}_+$ such that

$$\varphi_1(d(x,y)) \le d(f(x), f(y)) \le \varphi_2(d(x,y))$$

for every $x, y \in X$.

Since X is a geodesic space, by Lemma 2.1 one may choose φ_2 linear, $\varphi_2(t) = at + b$, $a > 0$. There exists $m_0 \in \mathbb{N}$ such that $d(f(x), f(y)) \ge 1$ as $d(x,y) \ge m_0$.

For any $m \ge m_0$, find minimal $k(m) \in \mathbb{N}$ such that the preimage $f^{-1}(\mathcal{A})$ of any $d_{k(m)}$-discrete family is $3m$-discrete. We have for $x \in A$, $y \in B$, $A, B \in \mathcal{A}$, $A \ne B$:

$$d_{k(m)} \le d(f(x), f(y)) \le \varphi_2(d(x,y)) = ad(x,y) + b$$

whence $d(x,y) \ge (d_{k(m)} - b)/a$ and if $(d_{k(m)} - b)/a \ge 3m$, we are done. Therefore, the minimal $k(m)$ is found from the inequality $d_{k(m)} \ge 3ma + b$.

We see that there exists $m_1 \in \mathbb{N}$ such that $d_m \ge (3m - b)/a$ for every $m \ge m_1$.

Now consider $m > \max\{m_0, m_1\}$. The cover $\mathcal{V}_m$ splits into the union of d_m-disjoint families $\mathcal{V}_m^i$, $i = 0, 1 \dots, n$.

The cover $f^{-1}(\mathcal{V}_m)$ of $X(m, R_m)$ splits into the union of $3m$-disjoint families $f^{-1}(\mathcal{V}_m^i)$, $i = 0, \dots, n$. By Proposition 3.14, there exists an element $V \in \mathcal{V}_m$ such that $K_m(f^{-1}(V)) \ge \sum_{i=1}^m c_i + 1 > c_m$. As $m > m_0$, we see that $K_1(V) > c_m$, a contradiction.

□

4. HIGSON PROPERTY

A metric space X with $\operatorname{asdim} X \le n$ is said to satisfy the *Higson property* if there exists $C > 0$ such that for every $D > 0$ there exists a cover $\mathcal{U}$ of X with $\operatorname{mesh}(\mathcal{U}) < CD$ and such that $\mathcal{U} = \mathcal{U}^0 \cup \dots \cup \mathcal{U}^n$, where $\mathcal{U}^0, \dots, \mathcal{U}^n$ are D-disjoint. Equivalently, X has the Higson property if there exists a sequence $(\mathcal{U}_k)$ of uniformly bounded covers of X and $C > 0$ such that $\operatorname{mesh}(\mathcal{U}_k) \to \infty$ and every ball in X of radius $\le C \operatorname{mesh}(\mathcal{U}_k)$ intersects at most $n+1$ element of $(\mathcal{U}_k)$. The latter is a large scale analog of the *Nagata dimension* N-dim defined by Assouad [1] as follows: for a metric space X we have $\text{N-dim}\, X \le n$ if there is a constant C such that, for each $r > 0$, there is an open cover $\mathcal{U}(r)$ of X by sets of diameter $\le Cr$ such that each open ball of radius r meets at most $n+1$ members of $\mathcal{U}(r)$.

Proposition 4.1. *Every proper metric space of finite asymptotic dimension is coarsely equivalent to a proper metric space that satisfies the Higson property.*

Proof. By Theorem 3.5, there is a large scale embedding of X into a space of the form $M(T^0,\dots,T^n)$, for a sequence of regular locally finite trees $T^0,\dots,T^n$ together with filtrations $T^i_0 \supset T^i_1 \supset T^i_2 \supset \dots$. Since the Higson property is preserved by subspaces, it is sufficient to show that the space $M(T^0,\dots,T^n)$ satisfies it.

Note that the set $K_j = r_j(M(T^0,\dots,T^n)$ is a cubic piecewise euclidean n-dimensional complex with $\operatorname{mesh}(K_j) = D_j = 2^j$. Passing to the second barycentric subdivision of K_j we are able to produce $D_k/3$-discrete families $\mathcal{A}_l$ with $\operatorname{mesh}(\mathcal{A}_l) \le 2D_k/3 \le D_k$, $l \in \{0,\dots,n\}$. Let $\mathcal{U}^l_k = \{r_k^{-1}(A) \mid A \in \mathcal{A}_l\}$, then

$$\operatorname{mesh}(\mathcal{U}^l_k) \le 2(D_k + D_{k-1} + \dots + D_0) \le 4D_k.$$

Since the map r_k is short, the family $\mathcal{U}^l_k$ is also $D_k/3$-discrete. □

A metric space X satisfies the *n-dimensional Nagata property* (briefly denoted by (P_n)) if for every $r > 0$, every $x \in X$, and every $y_1,\dots,y_{n+2}$ such that $d(y_i, N_r(x)) < 2r$ for every $i = 1,\dots,n+2$, there exist $i,j \in \{1,\dots,n+2\}$, $i \ne j$, such that $d(y_i,y_j) < 2r$ (see [15]).

Theorem 4.2. *For a proper metric space X the following are equivalent:*

(1) $\operatorname{asdim} X \le n$;
(2) *X is large scale equivalent to a metric space with property* (P_n).

Proof. (1)$\Longrightarrow$(2) By Proposition 4.1, X is coarsely equivalent to a proper metric space with the Higson property. Moreover, as we can see from the proof of Proposition 4.1, without loss of generality, we may assume that there exists a sequence $(\mathcal{U}_k)$ of uniformly bounded open covers of X and $C > 0$ such that $\operatorname{mesh}(\mathcal{U}_k) = 2^k$ and $\mathcal{U}_k$ splits into the union of $n{+}1$ families that are $C2^k$-discrete. Equivalently, every ball of radius $C2^{k-1}$ intersects at most $n+1$ element of the cover $\mathcal{U}_k$. By Lemma 2.2, there exists a 1-discrete subset X' of X which is coarsely equivalent to X. For every $r \in (0,\infty)$, put

$$\mathcal{U}'(r) = \begin{cases} \{\{x\} \mid x \in X'\} & \text{whenever } 0 < r < 1, \\ \{U \cap X' \mid U \in \mathcal{U}_{[r]}\} & \text{whenever } r \ge 1, \end{cases}$$

and $C' = C/4$. Then every ball in X' of radius $\le C'r$ intersects at most $n+1$ element of the cover $\mathcal{U}'(r)$, $r \in (0,\infty)$. The latter means that $\text{N-dim}\, X' \le n$ and by

It is proved in [1][Proposition 2.2] that then there exists a metric δ on X' such that δ satisfies $(\mathrm{P})_n$ and δ^p is Lipschitz equivalent to the original metric,

d, on X'. The latter means that there exist $A, B > 0$ such that

$$A\delta(x,y)^p \le d(x,y) \le B\delta(x,y)^p,\ x,y \in X'.$$

(2)$\Longrightarrow$(1) The proof is completely parallel to that in the case of dimension dim (see [15]). Let X be a proper metric space whose metric, d, satisfies property (P_n). Given $D > 0$, find a maximal (with respect to inclusion) set $Y \subset X$ which is $2D$-discrete. Then $\mathcal{U} = \{N_{2D}(y) \mid y \in Y\}$ is a uniformly bounded cover of X.

Show that for every $x \in X$ the set $N_D(x)$ intersects at most $n+1$ element of $\mathcal{U}$. Indeed, suppose the contrary; then there $y_1, \dots, y_{n+2} \in Y$ such that $N_D(x) \cap N_{2D}(y_i) \ne \emptyset$ for every $i = 1, \dots, n+2$. Property (P_n) implies that there exist $i, j \in \{1, \dots, n+2\}$, $i \ne j$, such that $d(y_i, y_j) < 2D$. This is a contradiction with the choice of Y. Because of arbitrarity of D, $\operatorname{asdim} X \le n$.

□

5. Coincidence of asymptotic dimensions

5.1. **Higson compactification and Higson corona.** Let $\varphi\colon X \to \mathbb{R}$ be a function defined on a metric space X. For every $x \in X$ and every $r > 0$ let $V_r(x) = \sup\{|\varphi(y) - \varphi(x)| \mid y \in N_r(x)\}$. A function φ is called *slowly oscillating* whenever for every $r > 0$ we have $V_r(x) \to 0$ as $x \to \infty$ (the latter means that for every $\varepsilon > 0$ there exists a compact subspace $K \subset X$ such that $|V_r(x)| < \varepsilon$ for all $x \in X \setminus K$. Let $\bar{X}$ be the compactification of X that corresponds to the family of all continuous bounded slowly oscillation functions. The *Higson corona* of X is the remainder $\nu X = \bar{X} \setminus X$ of this compactification.

It is known that the Higson corona is a functor from the category of proper metric space and coarse maps into the category of compact Hausdorff spaces. In particular, if $X \subset Y$, then $\nu X \subset \nu Y$.

For any subset A of X we denote by A' its trace on νX, i. e. the intersection of the closure of A in $\bar{X}$ with νX. Obviously, the set A' coincides with the Higson corona νA.

5.2. **Asymptotic inductive dimensions.** The notion of asymptotic inductive dimension asInd is introduced in [10].

Recall (see, e.g. [12]) that a closed subset C of a topological space X is a *separator* between disjoint subsets $A, B \subset X$ if $X \setminus C = U \cup V$, where U, V are open subsets in X, $U \cap V = \emptyset$, $A \subset U$, $V \subset B$.

Let X be a proper metric space. A subset $W \subset X$ is called an *asymptotic neighborhood* of a subset $A \subset X$ if $\lim_{r\to\infty} d(X \setminus N_r(x_0), X \setminus W) = \infty$. Two sets A, B in a metric space are *asymptotically disjoint* if $\lim_{r\to\infty} d(A \setminus N_r(x_0), B \setminus N_r(x_0)) = \infty$. In other words, two sets are asymptotically disjoint if the traces A', B' on νX are disjoint.

A subset C of a metric space X is an *asymptotic separator* between asymptotically disjoint subsets $A_1, A_2 \subset X$ if the trace C' is a separator in νX between A'_1 and A'_2.

By the definition, $\operatorname{asInd} X = -1$ if and only if X is bounded. Suppose we have defined the class of all proper metric spaces Y with $\operatorname{asInd} Y \leq n-1$. Then $\operatorname{asInd} X \leq n$ if and only if for every asymptotically disjoint subsets $A_1, A_2 \subset X$ there exists an asymptotic separator C between A_1 and A_2 with $\operatorname{asInd} C \leq n-1$. The dimension function asInd is called the *asymptotic inductive dimension*.

We can similarly define the small inductive asymptotic dimension asind. By the definition, $\operatorname{asind} X = -1$ if and only if X is bounded. Suppose we have defined the class of all proper metric spaces Y with $\operatorname{asind} Y \leq n-1$. Then $\operatorname{asind} X \leq n$ if and only if for every $A \subset X$ and every $x \in \nu X \setminus A'$ there exists a subset C of X such that C' is a separator between A' and $\{x\}$ in νX and $\operatorname{asind} C \leq n-1$.

Proposition 5.1. *For every metric space X we have* $\operatorname{asind} X \leq \operatorname{asInd} X$.

Proof. Induction by $\operatorname{asInd} X$. By the definition, the properties $\operatorname{asind} X = -1$ and $\operatorname{asInd} X = -1$ are equivalent. Suppose $\operatorname{asInd} X \leq n$ and $A \subset X$, $x \in \nu X \setminus A'$. There exists a continuous function $\varphi\colon \bar{X} \to [-1,1]$ such that $\varphi|\bar{A} = -1$, $\varphi(x) = 1$. The sets A and $B = (\varphi|X)^{-1}([0,1]) = \varphi^{-1}([0,1]) \cap X$ are asymptotically disjoint subsets of X ad therefor there exists an asymptotic separator C between A and B with $\operatorname{asInd} C \leq n-1$. Then obviously, C' is a separator between A' and x in νX and, by inductive hypothesis, $\operatorname{asind} C \leq n-1$. Therefore $\operatorname{asind} X \leq n$. □

The following is a counterpart of Theorem 1 in [10].

Proposition 5.2. *Let X be a proper metric space. Then* $\operatorname{asind} X \geq \operatorname{ind} \nu X$.

Proof. Induction by $\operatorname{asind} X$. Obviously, $\operatorname{asind} X = -1$ if and only if $\operatorname{ind}\nu X = -1$. Suppose that $\operatorname{asind} X \leq n$ and show that $\operatorname{ind}\nu X \leq n$. Let A be a closed subset of νX and $x \in \nu X \setminus A$. There exists a continuous function $\varphi\colon \bar{X} \to [-1,1]$ such that $\varphi|A = -1$, $\varphi(x) = 1$. Let $B = \varphi^{-1}([-1,0])$. Then $x \notin B'$ and, since $\operatorname{asind} X \leq n$, there exists a subset $C \subset X$ with $\operatorname{asind} C \leq n-1$ and such that C' is a separator between B' and x in νX. By the inductive hypothesis,

$$\operatorname{ind} C' = \operatorname{ind}\nu C \leq \operatorname{asind} C \leq n-1.$$

Since C is a separator between B' and x in νX, it is also a separator between A and x.

Therefore, for every closed subsert A of νX and $x \in \nu X \setminus A$ we are able to find a separator K with $\operatorname{ind} K \leq n-1$. This means that $\operatorname{ind}\nu X \leq n$. □

It is proved in [10] (see Proposition 1 therein) that $\operatorname{asInd} X = \operatorname{asInd} Y$ for coarsely equivalent spaces X and Y.

Proposition 5.3. *Let X and Y be coarsely equivalent metric spaces. Then* $\operatorname{asind} X = \operatorname{asind} Y$*.*

Proof. Analogous to that of Proposition 1 in [10]. $\square$

Lemma 5.4. *Suppose that $A \cup B$ are asymptotically disjoint subsets of $X \subset Y$ and C is an asymptotic separator in Y between A and B with* $\operatorname{asdim} C \leq m$*. Then there exists an asymptotic separator $\tilde{C}$ between A and B in X with* $\operatorname{asdim} \tilde{C} \leq m$

Proof. Define subsets D_k of X by induction. Let $D_1 = N_1(C) \cap X$ and $D_{k+1} = (N_{k+1}(C) \cap X) \setminus N_k(D_k)$. By Lemma 2.2, there exists a $(k+1)$-net $\tilde{D}_{k+1}$ in D_{k+1} which is k-discrete. Let $\tilde{C} = \cup_{k=1}^{\infty} \tilde{D}_k$.

We first show that $\operatorname{asdim} \tilde{C} \leq m$. Remark that for every $k > 0$ the complement to the subset $\cup\{\tilde{D}_j \mid j \leq k\}$ is the union of a k-disjoint family $\{\tilde{D}_j \mid j > k\}$. To apply Theorem 2.5, we have to verify that $\operatorname{asdim} \tilde{D}_k \leq m$ uniformly. If $k > 0$, then for some $R > 0$ there exists an R-bounded cover $\mathcal{U}$ of the set $\cup\{\tilde{D}_j \mid j \leq k\}$ that splits into the union $\mathcal{U}^0 \cup \dots \cup \mathcal{U}^m$ of k-discrete families. Let $\mathcal{V}^0 = \mathcal{U}^0 \cup \{\{x\} \mid x \in \cup\{\tilde{D}_j \mid j > k\}\}$, then the family $\mathcal{V}^0 \cup \mathcal{U}^1 \cup \dots \cup \mathcal{U}^m$ is a $\min\{R, k\}$-uniformly bounded family that splits into the union of $m+1$ k-disjoint families.

Show that for every integer $k > 0$ there exists $R = R(k) > 0$ such that $N_k(C) \cap X \subset N_{R(k)}(\tilde{C})$. Indeed, if $x \in N_k(C) \cap X$, then either $x \in D_k \subset N_{2k}(\tilde{D}_k)$ or

$$x \in N_k(C) \setminus D_k \subset N_{k-1}(D_{k-1}) \subset N_{k-1}(N_{k-1}(\tilde{D}_{k-1})) \subset N_{2k-2}(\tilde{D}_{k-1}) \subset N_{2k-2}(\tilde{C})$$

and we can choose $R(k) = 2k$.

Now suppose that C is an asymptotic separator between A and B in Y. Show that $\tilde{C}$ is an asymptotic separator between A and B in X. It is sufficient to show that $\tilde{C}' \supset C' \cap \nu X$. Suppose that $a \in C' \cap \nu X$ and U is a closed neighborhood of a in $\bar{Y}$. Then there exist sequences (x_i) in $X \cap U$ and (c_i) in $C \cap U$ and $k \in \mathbb{N}$ such that $d(x_i, c_i) \leq k$ for every i.

Case 1). For infinite number of i (passing to a subsequense we then assume that for all i) $x_i \in D_k$. Then there exist $\tilde{x}_i \in \tilde{D}_k$ with $d(x_i, \tilde{x}_i) \leq k$.

Case 2). For infinite number of i (passing to a subsequense we then assume that for all i) $x_i \in N_{k-1}(D_{k-1})$. Then there exist $y_i \in D_{k-1}$ such that $d(x_i, y_i) \leq k-1$ and there exist $\tilde{x}_i \in \tilde{D}_{k-1}$ such that $d(y_i, \tilde{x}_i) \leq k-1$.

In both cases, $\tilde{x}_i \in \tilde{C}$ for every i and $d(x_i, \tilde{x}_i) \leq \max\{k, 2k-1\} = 2k$ and therefore $\tilde{x}_i \in U \cap X$ for all but finitely many i. This means that the sets $\tilde{C}$ and U are not asymptotically disjoint and therefore $C' \cap U \neq \emptyset$. Because of arbitrarity of U we conclude that $a \in \tilde{C}'$. $\square$

Theorem 5.5. *For every space $X \in \mathcal{M}_n$ we have* $\operatorname{asInd} X \leq n$*.*

We need some auxiliary results. Let $T^0, \dots, T^n$ be a sequence of regular locally finite trees.

Proposition 5.6. *For every disjoint asymptotically disjoint subsets A, B in $M(T^0, \dots, T^n)$ there exists a separator C between them such that* $\operatorname{asdim} C \leq n-1$ *and the sets A, B, and C are asymptotically disjoint.*

Proof. Let

$$\begin{aligned} U =& \{x \in M(T^0, \dots, T^n) \mid d(x, A) \leq (1/2)d(x, B)\}, \\ V =& \{x \in M(T^0, \dots, T^n) \mid d(x, B) \leq (1/2)d(x, A)\}, \end{aligned}$$

then U, V are closed, disjoint, asymptotically disjoint asymptotic neighborhoods of the sets A, B respectively. For every j let R_j be a number such that

$$d(r_j(U \setminus N_{R_j}(x_0)), r_j(V \setminus N_{R_j}(x_0))) \geq 2^{j+2}$$

(such a number exists because U, V are asymptotically disjoint and the retraction r_j is 2^{j+1}-close to the identity map. We will assume that $R_{j+1} \geq 2R_j$. Put $U_j = r_j\left(U \cap (\overline{N_{R_{j+1}}(x_0)} \setminus N_{R_j}(x_0))\right)$.

Denote by $\hat{U}_j$ the union of all n-dimensional cubic simplices in the subcomplex $r_j(M(T^0, \dots, T^n))$ that intersect U_j. Let $S_j = r^{-1}(\partial(N_{1/2}(\hat{U}_j))$ and $S = \cup_{i=0}^{\infty} S_j$.

We are going to show that S contains a separator between U and V and $\operatorname{asdim}(S) \leq n-1$.

Put $S' = \cup\{S_{2k} \mid k \in \mathbb{N}\}$, $S'' = \cup\{S_{2k-1} \mid k \in \mathbb{N}\}$. By finite sum theorem for asymptotic dimension (see, e.g. [2]), it is sufficient to prove that $\operatorname{asdim} S' \leq n-1$, $\operatorname{asdim} S'' \leq n-1$ and, because of complete analogy, we only prove that $\operatorname{asdim} S' \leq n-1$.

Prove that the family $\{S_{2k}\}$ has asymptotic dimension $\leq n-1$ uniformly. Let $D > 1$. There exists natural j such that every $(n-1)$-dimensional cubic complex of mesh 2^j can be covered by a 2^j-bounded family of its subsets that splits into the union of n families which are $2D$-disjoint.

Claim. $r_{jj'}^{-1}(\partial(N_{1/2}(\hat{U}_{j'})) \subset N_{3/4}(r_{jj'}^{-1}(\hat{U}_{j'}))^{(n-1)}$.

To prove Claim, suppose that $r_{jj'}(x) \in \partial(N_{1/2}(\hat{U}_{j'})$. Then there is a cubic n-dimensional symplex K in $\hat{U}_{j'}$ such that $r_{jj'}(x) \in \partial(N_{1/2}(K)$. The symplex K is of the form $K = \prod_{i=0}^{n} K_i$, where K_i is a 1-dimensional symplex in $T^i_{j'}$ and there is $l \in \{0, \dots, n\}$ such that K_l is a singleton, $K_l = \{a_l\}$. Then $\partial N_{1/2}(K) = (\cup_{k=0}^{n} R_k) \cap M_{j'}$, where

$$R_k = \left(\prod_{p \in \{0,\dots,n\} \setminus \{k\}} \overline{N_{1/2}(K_p)}\right) \times \partial N_{1/2}(K_k).$$

Then $r_{jj'}^{-1}(\partial N_{1/2}(K_k)) = \partial N_{1/2}(K_k)$ and we obtain

$$r_{jj'}^{-1}(R_k) \cap M_{j'} = \left(\prod_{p \in \{0,\dots,n\} \setminus \{k\}} (r_{jj'}^{p})^{-1}(\overline{N_{1/2}(K_p)}) \right) \times \partial N_{1/2}(K_k) \cap M_{j'}.$$

For every $x = (x_0, \dots, x_n) \in r_{jj'}^{-1}(R_k) \cap M_{j'}$ there exists $p \in \{0, \dots, n\} \setminus \{k\}$ such that $x_p \in \partial(T_{j-1}^{p} \setminus T_j^{p})$. Then $d(x, (x_1, \dots, x_{k-1}, y_k, x_{k+1}, \dots, x_n)) = 1/2$ for some $y_k \in \partial K_k$ (the boundary is considered in T^k; note that then $y_k \in r_{j'}^{k}(\partial T_{j'-1}^{k} \setminus T_{j'}^{k}))$ and $(x_1, \dots, x_{l-1}, a_l, x_{l+1}, \dots, x_n) \in N_{3/4}(r_{jj'}^{-1}(\hat{U}_{j'}))^{(n-1)}$. Claim is proved.

Denote by $\mathcal{V}$ a cover of the cubic symplex M_j that splits into the union $\mathcal{V} = \mathcal{V}^0 \cup \dots \cup \mathcal{V}^n$ of uniformly 2^j-bounded covers which are $2D$-discrete. Now for every k and every $i \in \{0, \dots, n\}$ consider the family $\tilde{\mathcal{V}}_k^i = \{N_{3/4}(V) \cap r_{j,2k}^{-1}(\hat{U}_{2k}) \mid V \in \mathcal{V}_k^i\}$ of the set $r_{j,2k}^{-1}(\hat{U}_{2k})$. The cover $\mathcal{V}_k = \mathcal{V}_k^0 \cup \dots \cup \mathcal{V}_k^n$ is uniformly $(2^j + 3)$-bounded and splits into the union of D-discrete families $\tilde{\mathcal{V}}_k^i$, $i \in \{0, \dots, n\}$. Then the families $\mathcal{W}_k^i = \{r_j^{-1}(V) \in V \in \mathcal{V}_k^i\}$ are uniformly $(2^{j+1} + 3)$-bounded and D-discrete and, because $S_{2k} = r_j^{-1} r_{j,2k}^{-1}(\hat{U}_{2k})$, we see that $\cup_{i=0}^{n} \mathcal{W}_k^i$ covers S_{2k}.

Applying Theorem 2.5 we conclude that $\operatorname{asdim} S \leq n - 1$. Finally, note that S contains a separator between A and B, namely, the set $\partial N_{1/2}(\cup_{j=1}^{\infty} \hat{U}_j)$. $\square$

Proposition 5.7. *Let C be a separator between asymptotically disjoint subsets A, B in a proper uniformly arcwise connected space X and the sets C and $A \cup B$ are asymptotically disjoint. Then C is an asymptotic separator between A and B in X.*

Proof. Denote by U the set of points that can be connected to A by an arc in the set $X \setminus C$. Show that the sets $(\bar{U}) \cap X$ and B are asymptotically disjoint. Indeed, assuming the opposite we obtain sequences (a_i) and (b_i) in U and B respectively such $d(a_i, x_0) \to \infty$, $d(b_i, x_0) \to \infty$ (here x_0 is an arbitrary base poin in X) and that there exists $R > 0$ for which $d(a_i, b_i) < R$, $i \in \mathbb{N}$. Since X is uniformly arcwise connected, there exists $S > 0$ such that, for every i, the points a_i and b_i can be connected by an arc of diameter $\leq S$. There exists natural i_0 such that for every $i \geq i_0$ we have $d(b_i, C) \geq R + S$. Then obviously there exists an arc in $X \setminus C$ that connects a_i and b_i, $i \geq i_0$. Therefore, $b_i \in U$ and we obtain a contradiction.

Thus, $B' \subset (X \setminus U)'$. In order to prove that C is an asymptotic separator between A and B it remains to prove that $\bar{U} \cap \bar{B} \cap \nu X \subset C'$. Indeed, suppose the opposite, i.e. that there exists a point $y \in \bar{U} \cap \bar{B} \cap \nu X \setminus C'$. Then there exist disjoint neighborhoods V of y and W of $\bar{C}$ in $\bar{X}$ and obviously $\overline{U \cap V} \cap \bar{B} \neq \emptyset$. Therefore, there exist sequences (a_i) in $U \cap V$ and (b_i) in B such that $\lim_{i \to \infty} d(a_i, x_0) = \infty$, $\lim_{i \to \infty} d(b_i, x_0) = \infty$, and the sequence $(d(a_i, b_i))$ is bounded. Arguing similarly as above we conclude that then there exists a sequence (c_i) in C for which the sequence $(d(a_i, c_i))$ is bounded. This contradicts to our choice of V.

□

Lemma 5.8. *The space $M(T^0, \dots, T^n)$ is uniformly arcwise connected.*

Proof. Note that for every l and every $z \in r_l(M(T^0, \dots, T^n)) \setminus r_{l+1}(M(T^0, \dots, T^n))$ there exists a path of diameter $\leq 2^{l+1}$ in $M(T^0, \dots, T^n)$ connecting z with $r_l(z)$.

Let $x = (x_0, \dots, x_n), y = (y_0, \dots, y_n) \in M(T^0, \dots, T^n)$. Let j be the minimal natural number such that $d(x, y) < 2^j$. Then $r_j(x) = r_j(y)$. Let J be a path connecting x and y that consists of paths of diameter $\leq 2^{l+1}$ connecting $r_{l-1}(x)$ with $r_l(x)$ and of paths of diameter $\leq 2^{l+1}$ connecting $r_{l-1}(y)$ with $r_l(y)$, $1 \leq l \leq j$ (recall that r_0 is the identity map). Then

$$\begin{aligned}\operatorname{diam}(J) \leq & \sum_{l=1}^{j} d(r_{l-1}(x), r_l(x)) + \sum_{l=1}^{j} d(r_{l-1}(y), r_l(y)) \\ \leq & 2\sum_{l=1}^{j} 2^{l+1} \leq 2^{j+3} \leq 16 d(x, y).\end{aligned}$$

□

Now we are going to prove Theorem 5.5. It is proved in [10] that $\operatorname{asInd} X \geq \operatorname{Ind} \nu X$. This result together with the classical theorem on the comparison of Ind and dim implies the inequality $\operatorname{asInd} X \geq \dim \nu X$. Suppose $\operatorname{asdim} X < \infty$. Then $\operatorname{asdim} X = \dim \nu X$ (see [7]), and we obtain $\operatorname{asInd} X \geq \operatorname{asdim} X$ for every X with $\operatorname{asdim} X < \infty$.

To prove the opposite inequality we apply induction on $\operatorname{asdim} X$. It is known (see [10]) that the properties $\operatorname{asdim} X = 0$ and $\operatorname{asInd} X = 0$ are equivalent. Assume that the inequality $\operatorname{asInd} X \leq \operatorname{asdim} X$ is proved for every X with $\operatorname{asdim} X \leq n - 1$.

Suppose that $\operatorname{asdim} X \leq n$. Then by Theorem 3.5 there is a large scale embedding of X into the space $M(T^0, \dots, T^n)$, for some regular locally finite trees $T^0, \dots, T^n$. Since the image of any space under a large scale embedding is coarse equivalent to this space and the dimension asInd is a coarse invariant (see [10, Proposition 1]), we may assume that X is a subspace of $M(T^0, \dots, T^n)$. Let A, B be asymptotically disjoint subsets of X. Without loss of generality we assume A, B to be closed and disjoint. By Proposition 5.6, there exists a separator C between A and B with $\operatorname{asdim} C \leq n - 1$. Since, by Lemma 5.8, the space $M(T^0, \dots, T^n)$ is uniformly arcwise connected, it follows from Proposition 5.7 that C is also an asymptotic separator between A and B in $M(T^0, \dots, T^n)$.

By Lemma 5.4, there exists an asymptotic separator $\tilde{C}$ between A and B in X with $\operatorname{asdim}\tilde{C} \leq n-1$. Applying the induction assumption we see that $\operatorname{asInd}\tilde{C} \leq n-1$.

Since for every asymptotically disjoint subsets in X there exists an asymptotic separator between them whose dimension asInd does not exceed $n-1$, we conclude that $\operatorname{asInd} X \leq n$.

The equality $\operatorname{asInd} X = \operatorname{asdim} X$ is therefore proven for all X with $\operatorname{asdim} X < \infty$.

One can similarly prove the following result.

Theorem 5.9. *For all proper metric spaces X with $\operatorname{asdim} X < \infty$ we have $\operatorname{asind} X = \operatorname{asdim} X$.*

We finally obtain that the dimensions asdim, asInd, and asind coincide in the class of proper metric spaces that are finite dimensional with respect to asdim. This fact can be regarded as a counterpart in the asymptotic topology of the classical result on coincidence of the covering dimension, the small inductive dimension, and the large inductive dimension (see [12]).

6. Open problems

The coincidence of the dimension functions asdim and asInd is proved under the assumption of finiteness of asdim. This leads to the following natural question.

Question 6.1. Is there a proper metric space X with $\operatorname{asdim} X = \infty$ and $\operatorname{asInd} X < \infty$?

A similar question can be formulated for ind.

A closed subset C of a topological space X is a *cut* between disjoint subsets $A, B \subset X$ if every continuum (compact connected space) $T \subset X$ that intersect both A and B also intersects C.

The notion of the large inductive dimension can be based on the notion of cut instead of separator. It turnes out (see [13]) that the obtained dimension (it was defined by Brouwer who called it Dimensiongrad) coincides with the classical large inductive dimension in the class of separable metrizable spaces).

Question 6.2. Does the large asymptotic inductive dimension defined on the base of asymptotic cut coincides with asInd?

References

[1] P. Assouad, *Sur la distance de Nagata.* C. R. Acad. Sci. Paris Sér. I Math. 294 (1982), no. 1, 31–34.

[2] G. Bell and A. Dranishnikov, *On asymptotic dimension of groups*, Algebraic and Geometric Topology 1 (2001), 57-71.

[3] G. Bell and A. Dranishnikov, *On asymptotic dimension of groups acting on trees*, Preprint.

[4] G. Bell, A. Dranishnikov and J. Keesling, *On a formula for the asymptotic dimension of amalgamated products,* Preprint.

[5] M. Bestvina, Characterizing k-dimensional universal Menger compacta. 71 (1988), no. 380, vi+110 pp.

[6] Ph. L. Bowers *General position properties satisfied by finite products of dendrites.* Trans. Amer. Math. Soc. 288 (1985), no. 2, 739–753.

[7] A. N. Dranishnikov, J. Keesling, V. V. Uspenskij, *On the Higson corona of uniformly contractible spaces.* Topology 37 (1998), no. 4, 791–803.

[8] A. Dranishnikov, *Asymptotic topology,* Russian Math. Surveys **55**(2000), No 6, 71–116.

[9] A. Dranishnikov, *On hypersphericity of manifolds with finite asymptotic dimension,* Preprint.

[10] A. Dranishnikov, *On asymptotic inductive dimension,* JP J. Geom. Topol. 1 (2001), no. 3, 239–247.

[11] A. Dranishnikov and T. Januszkiewicz, *Every Coxeter group acts amenably on a compact space.* Proceedings of the 1999 Topology and Dynamics Conference (Salt Lake City, UT). Topology Proc. 24 (1999), Spring, 135–141.

[12] R. Engelking, Theory of dimensions, finite et infinite. – Lemgo: Heldermann, 1995. (Sigma series in pure mathematics; Vol. 10).

[13] V. V. Fedorchuk, M. Levin, E. V. Shchepin, *On the Brouwer definition of dimension. (Russian)* Uspekhi Mat. Nauk 54 (1999), no. 2(326), 193–194; translation in Russian Math. Surveys 54 (1999), no. 2, 432–433.

[14] M. Gromov, *Asymptotic invariants for infinite groups,* LMS Lecture Notes, **182**(2), 1993.

[15] J. Nagata, *Note on dimension theory for metric spaces.* Fund. Math. 45 1958 143–181.

[16] Ph. A. Ostrand, *Dimension of metric spaces and Hilbert's problem* 13. Bull. Amer. Math. Soc. **71**(1965), 619–622.

[17] J. Roe, *Coarse cohomology and index theory for complete Riemannian manifolds,* Memoirs Amer. Math. Soc. No 497, 1993.

[18] E. V. Shchepin, *Arithmetic of dimension theory. (Russian)* Uspekhi Mat. Nauk 53 (1998), no. 5(323), 115–212; translation in Russian Math. Surveys 53 (1998), no. 5, 975–1069.

[19] G. Yu, *The Novikov conjecture for groups with finite asymptotic dimension.* Ann. of Math. (2) 147 (1998), no. 2, 325–355.

Department of Mathematics, University of Florida, 358 Little Hall, PO Box 118105, Gainesville, FL 32611–8105

E-mail address: `dranish@math.ufl.edu`

Department of Mechanics and Mathematics, Lviv National University, Universytetska 1, 79000 Lviv, Ukraine

E-mail address: `topos@franko.lviv.ua, mzar@litech.lviv.ua`